



Aleve® - D Sinus & Headache Caplets (NSAID Labeling)

Drug Facts	
Active ingredients (in each caplet)	Purposes
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*.....	Pain reliever/fever reducer
Pseudoephedrine HCl 120 mg, extended-release.....	Nasal decongestant *nonsteroidal anti-inflammatory drug
<p>Uses temporarily relieves these cold, sinus, and flu symptoms:</p> <ul style="list-style-type: none"> • sinus pressure • minor body aches and pains • headache • nasal and sinus congestion (promotes sinus drainage and restores freer breathing through the nose) • fever 	
<p>Warnings</p> <p>Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:</p> <ul style="list-style-type: none"> • hives • facial swelling • asthma (wheezing) • shock • skin reddening • rash • blisters <p>If an allergic reaction occurs, stop use and seek medical help right away.</p> <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:</p> <ul style="list-style-type: none"> • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed 	
<p>Do not use</p> <ul style="list-style-type: none"> • if you have ever had an allergic reaction to any other pain reliever/fever reducer • right before or after heart surgery • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. • in children under 12 years of age 	
<p>Ask a doctor before use if</p> <ul style="list-style-type: none"> • the stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease • you are taking a diuretic • you have problems or serious side effects from taking pain relievers or fever reducers • you have <ul style="list-style-type: none"> • asthma • diabetes • thyroid disease • trouble urinating due to an enlarged prostate gland 	
<p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> • under a doctor's care for any serious condition • taking any other drug 	
<p>When using this product</p> <ul style="list-style-type: none"> • take with food or milk if stomach upset occurs • the risk of heart attack or stroke may increase if you use more than directed or for longer than directed 	



Aleve® - D Sinus & Headache Caplets (NSAID Labeling)

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- redness or swelling is present in the painful area
- any new symptoms appear
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing or the caplet feels stuck in your throat
- you get nervous, dizzy, or sleepless
- nasal congestion lasts more than 7 days

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- **swallow whole**; do not crush or chew
- **drink a full glass of water with each dose**
- adults and children 12 years and older: **1 caplet every 12 hours**; do not take more than 2 caplets in 24 hours
- children under 12 years: do not use

Other information

- **each caplet contains:** sodium 20 mg
- store at 20-25°C (68-77°F)
- store in a dry place

Inactive ingredients colloidal silicon dioxide, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, talc, titanium dioxide

Questions or comments? 1-800-986-0369 (Mon – Fri 9AM – 5PM EST) or www.AleveD.com